



PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁵ : A61K 37/02, 39/00, C07K 7/08 C07K 7/10, 13/00, 15/00 C07K 15/28	A1	(11) International Publication Number: WO 91/04745 (43) International Publication Date: 18 April 1991 (18.04.91)
(21) International Application Number: PCT/US90/05105 (22) International Filing Date: 13 September 1990 (13.09.90) (30) Priority data: 413,332 27 September 1989 (27.09.89) US 571,267 23 August 1990 (23.08.90) US (71) Applicant: ATHENA NEUROSCIENCES, INC. [US/ US]; 800F Gateway Boulevard, South San Francisco, CA 94080 (US). (72) Inventors: RUBIN, Lee, L. ; 122 Funston Avenue, San Francisco, CA 94118 (US). LIAW, Chen, W. ; 80 Maria- na Court, Redwood City, CA 94062 (US). TOMASELLI, Kevin, J. ; 485 Vallejo Street, San Francisco, CA 94133 (US).		(74) Agents: BLECHER, Melvin et al.; 611 West Sixth Street, 34th Floor, Los Angeles, CA 90017 (US). (81) Designated States: AT (European patent), BE (European patent), CA, CH (European patent), DE (European pa- tent)*, DK (European patent), ES (European patent), FR (European patent), GB (European patent), IT (Euro- pean patent), JP, LU (European patent), NL (European patent), SE (European patent). Published <i>With international search report.</i> <i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>

(54) Title: COMPOSITIONS FOR CELL ADHESION INHIBITION AND METHODS OF USE

(57) Abstract

Compositions that disrupt microvascular endothelial and epithelial cell tight junctions, and methods of use, are disclosed. Such compositions comprise agents that inhibit the binding to such cells of cell adhesion molecules. Such inhibitor agents include cell adhesion molecules, fragments of cell adhesion molecules that encompass a cell-binding domain such as HAV, and antibodies directed against cell adhesion molecules and fragments thereof. Also disclosed are drug delivery compositions comprising a therapeutic drug conjugated to an agent that disrupts cell tight junctions.

BEST AVAILABLE COPY

DESIGNATIONS OF "DE"

Until further notice, any designation of "DE" in any international application whose international filing date is prior to October 3, 1990, shall have effect in the territory of the Federal Republic of Germany with the exception of the territory of the former German Democratic Republic.

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	ES	Spain	MC	Monaco
AU	Australia	FI	Finland	MG	Madagascar
BB	Barbados	FR	France	ML	Mali
BE	Belgium	GA	Gabon	MR	Mauritania
BF	Burkina Fasso	GB	United Kingdom	MW	Malawi
BG	Bulgaria	GR	Greece	NL	Netherlands
BJ	Benin	HU	Hungary	NO	Norway
BR	Brazil	IT	Italy	PL	Poland
CA	Canada	JP	Japan	RO	Romania
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	LI	Liechtenstein	SN	Senegal
CM	Cameroon	LK	Sri Lanka	SU	Soviet Union
DE	Germany	LU	Luxembourg	TD	Chad
DK	Denmark			TC	Togo
				US	United States of America

THE PATENT OFFICE

**COMPOSITIONS FOR CELL ADHESION INHIBITION
AND METHODS OF USE**

This is a continuation-in-part of United States
Serial No. 07/413,332, filed September 27, 1989.

5 **Background of the Invention**

Field of the Invention

 This invention relates to compositions that
transiently and reversibly dissociate the blood-brain
barrier. More particularly, the invention relates to
10 compositions that dissociate tight junctions between
brain capillary endothelial cells that constitute the
physiological barrier between the general circulation
and the brain.

Detailed Description of Related Art

15 The entry of drugs from the blood stream to the
central nervous system (CNS), i.e., the brain and
spinal cord, is restricted by the presence of high
resistance tight junctions between brain capillary
cells and by the apparently low rate of transport
20 across these endothelial cells (Betz, A.L., et al.,
Ann. Rev. Physiol., 48:241 (1986); Pardridge, W.M.,
Ann. Rev. Pharmacol. Toxicol., 28:25 (1988)).

 The tight junctions of the blood brain barrier
(BBB) prevent diffusion of molecules and ions around
25 the brain capillary endothelial cells. The only
substances that can readily pass from the luminal core
of the capillary to the abluminal tissues that surround
the capillary are those molecules for which selective
transport systems exist in the endothelial cells, as
30 well as those compounds that are lipophilic (i.e.,
hydrophobic). In contrast, drugs, peptides and other

molecules that are neither lipophilic nor transported by specific carrier proteins are barred from entry into the brain, or their rates of entry are too low to be useful, thereby imposing a severe limitation upon the physician's ability to treat CNS disorders pharmacologically.

The carrier-mediated transcellular transport system mentioned above may have limited usefulness for therapeutic modalities under some circumstances.

Transcytotic transport, in general, involves, first, the binding of molecules to specific carrier proteins on the surface of endothelial cells, and, second, the delivery of such molecules across the endothelial cells. Limitations on the usefulness of such a system for treatment of CNS disorders are based on the following considerations: (1) physiological carrier proteins may not function efficiently, or at all, with non-physiological drugs; (2) even where function occurs, the rate of transport of therapeutic agents will be limited by the rate of transport of the carrier; (3) the overall capacity of cerebral capillary endothelial cells to transport any therapeutic macromolecules may be simply too low to achieve therapeutic levels of certain drugs in the brain; and (4) once therapeutic macromolecules enter endothelial cells, depending on their nature, they might be delivered to any number of organelles, including lysosomes that contain a wide variety of hydrolytic enzymes. For these reasons, creating drug delivery systems that do not rely upon transcytosis will clearly be advantageous.

As tight junctions between brain capillary endothelial cells constitute a major part of the BBB, the possibility of modifying these junctions has been considered. It has been found that tight junctions,

including those of the BBB, can be disrupted by hyperosmotic solutions administered intra-arterially. For example, Polley et al., WO89/04663, published June 1, 1989, disclose the osmotic disruption of the interendothelial structure of the BBB by the intra-arterial administration of hypertonic solutions of mannitol, arabinose or glycerol as a means of introducing into the brain genetic material. Similarly, hyperosmotic solutions of urea have also been used to alter the BBB (Bowman, P.D. et al., Ped. Res., 16:335A (1982)).

Other chemical agents have been reported to disrupt endothelial or epithelial cell tight junctions when administered intravenously, including:

7-fluorouracil (MacDonell, L.A., et al., Cancer. Res., 38:2930 (1978)), degradation by membrane enzymes (Vincent, P.A., et al., Exp. Mol. Path., 48:403 (1988); Diener, H.M., et al., J. Immunol., 135:537 (1985)), aluminum salts (Zigler, Z.Y., et al., IRCS Med. Sci., 12:1095 (1984)), histamine (Meyrick, B., et al., Exp. Lung Res., 6:11 (1984)), thrombin (Siflinger-Birnboim, A., et al., Microvasc. Res., 36:216 (1988)), phorbol esters (Shiba, K., et al., Exp. Cell Res., 178:233 (1988)), and neutralization of the luminal anionic charge (Hart, M.M., J. Neuropathol. Exp. Neurol., 46:141 (1987)).

Although the above-listed modalities may disrupt tight junctions and thereby increase permeability of the BBB, problems attendant upon their use make them less than desirable. For example, intra-arterial perfusion with hyperosmotic solutions involves surgery, and this cannot be repeated on a regular basis. Further, concentrated sugar solutions may not be innocuous, and might be expected to have undesirable side effects. In addition, the aforementioned chemical

agents may not be useful for the treatment of chronic neurological disease, their effects on tight junctions are not always reversible, and, as they all are themselves powerful drugs, there is always the danger that their use will compromise the patient's health generally. For example, 7-fluorouracil is a powerful inhibitor of pyrimidine synthesis, and thus nucleic acid biosynthesis, in animals cells.

Thus, an important need still exists for means which transiently and reversibly disrupt tight junctions of the BBB in order that administered drugs can reach the brain from the general circulation, and which have no undesirable side effects of their own in the subject.

Attempts have been made to disrupt cell-cell adhesion by modifying the protein(s) responsible for such adhesion, collectively referred to as "cell adhesion molecules" (CAM). One class of CAM is termed "cadherin". "Cadherin" is the term applied to a family of glycoproteins found in most kinds of mammalian tissues and thought to be responsible for Ca^{2+} -dependent cell-cell adhesion, (Takeichi, M., Development, 102:639 (1988)). Three subclasses of cadherin have been identified, namely, E-cadherin (from epithelial tissues), P-cadherin (from placental tissues), and N-cadherin (from neural tissues) (Yoshida-Noro, C., et al. Dev. Biol., 101:19 (1984); Nose, A., et al., J. Cell Biol., 103:2649 (1986); Hatta, K., et al., Nature, 320:447 (1986)).

The different cadherins exhibit distinct tissue distribution patterns (Takeichi, U., (1988) above). E-cadherin, which was found to be distributed exclusively in epithelial cells of various tissues (Hatta, K., et al., Proc. Nat'l. Acad. Sci. (USA), 82:2789 (1985); Takeichi, 1988, above), appears to be

identical to uvomorulin (Hyafil, F., et al., Cell, 21:927 (1986)), chicken liver-cell adhesion molecule (L-CAM, Gallin, W.J., et al., Proc. Nat. Acad. Sci. (USA), 80:1038 (1983)), and cell-CAM 120/80 (Damsky,

5 C.H., et al., Cell, 34:455 (1983)) in terms of biochemical properties (Cunningham, B.A., et al., Proc. Nat. Acad. Sci. (USA), 81:5787 (1984)) and tissue distributions (Thiery, J.-P., et al., Dev. Biol., 102:61 (1984)).

10 N-cadherin, which is expressed in various neural tissues including astrocytes (Hatta, K., et al., Devel. Biol., 120:215 (1987); Matsunaga, M., et al., Nature, 334:62 (1988); Tomaselli, K.J., Neuron, 1:33 (1988)), shows 92% amino acid sequence homology between
15 mammalian and avian homologs, shows from 40 to 50% similarity to epithelial E-cadherin and to placental P-cadherin of the same species, but was immunologically not cross-reactive with other cadherins within the same animal (Miyatani, S., Science, 245:631 (1989)).

20 Placental P-cadherin has also been cloned, and the deduced amino acid sequence of this glycoprotein was found to exhibit about 58% homology with epithelial E-cadherin (Nose, A., et al., EMBO J., 12:3655 (1987)).

Subsequent to the September 27, 1989 filing of the
25 parent application, Heimark, et al. (Heimark, R.L., et al., J. Cell Biol., 110:1745 (1990) reported on the identification of a Ca^{2+} -dependent cell-cell adhesion molecule in aortic endothelial cells.

Although each of the aforelisted cadherins
30 displays unique immunological and tissue distribution specifications, all have features in common: (1) a requirement for Ca^{2+} for cell adhesion function; (2) protection by Ca^{2+} from proteolytic cleavage; (3) similar numbers of amino acids, i.e., from about 723 to
35 about 822; (4) similar masses, i.e., about 124 kdal.

for the glycoprotein; (5) substantial interspecies (50%-60%) overall sequence homology with interspecies homologies increasing to about 56% to 99% in the cytoplasmic region of the protein, suggesting that they constitute a gene family (Nose, A., 1987; Miysysni, D., et al., 1989); and (6) a common mechanism of action, namely, homophilic binding of cadherins on one cell to similar cadherins on the adjoining cell.

CAMs independent of Ca^{2+} are also known, for example, the 125K glycoprotein of Urushihara et al. (Urushihara, H., et al., Cell, 20:363 (1980)); N-CAM (Rutishauser, U., Nature, Lond., 310:549 (1984)); Ng-CAM (Grunet, M. et al., Proc. Nat'l. Acad. Sci. (USA), 81:7989 (1984)); L1 (Rathjien, F.G. et al., ***** J., 3:1 (1984)); G4 (Rathjien, F.G. et al., J. Cell Biol., 104:343 (1987)); and platelet glycoprotein PECAM-1 (CD 31) (Newman, P.J., Science, 247:1219 (1990)). Ca^{2+} -independent CAMs are known to exhibit certain properties of the Ca^{2+} -dependent CAMs. Thus, N-CAM and N-cadherin both promote retinal neurite outgrowth on astrocytes (Neugebauer, K.M., et al., J. Cell Biol., 107:1177 (1985)), and on Schwann cells (Bixby, J.L. et al., J. Cell Biol., 107:353 (1988)).

Monoclonal antibodies raised against epithelial E-type cadherins such as uvomorulin are known to disrupt the adhesion of several cell types, including embryo cells, cultured teratocarcinoma cells, hepatocytes, and MDCK kidney epithelial cells (Ogou, S.-I., et al., J. Cell Biol., 97:944 (1983); Yoshida-Noro, et al., (1984), above; Shirayoshi, Y., et al., Cell Struct. Funct., 11:285 (1986); Gallin, et al., (1983), above; Vestweber, D., et al., EMBO J., 4:3393 (1985); Johnson, M.H., et al., J. Embrol. Exp. Morphol., 93:239 (1986); Gumbiner, B., et al., J. Cell Biol., 102:457 (1986)).

However, prior to the present discoveries disclosed in the parent applications cadherins had not been found in brain capillary or other endothelial cells (see, Takeichi, et al. (1988), above). Further, 5 the CAMs of microvascular endothelial cells had not yet been identified, nor had such molecules been localized specifically to brain capillary endothelial cells. Thus, until the present invention no means were known for transiently and reversibly disrupting tight 10 junctions between microvascular endothelial cells, including those of the BBB, based upon an attack upon the CAM's of such cells that are responsible for tight junction formation and maintenance.

It has been hypothesized that the cadherins 15 contain a common cell adhesion recognition (CAR) sequence. The CAR sequences of several cell and substratum adhesion molecules are known. Martin, G.R., et al., Ann. Rev. Cell Biol., 3:57 (1987) ; Ruoslahti, E., et al., Science, 238:491 (1987). In general, CAR 20 sequences are composed of at least three amino acid residues. The most rigorously investigated CAR sequence is RGD which is found in laminin, fibronectin and other basement membrane components that are responsible for the binding of cells to the substratum.

25 Blaschuk, et al., in a paper to be published subsequent to the filing of the present application (Blaschuk, O., et al., J. Mol. Biol., in press, (1990)), disclose the presence of three potential cadherin CAR sequences in the first extracellular 30 domains of liver CAM, E-, P-, and N-cadherin, namely, PPI, GAD and HAV. Blaschuk, et al. (Blaschuk, O., et al., Develop. Biol., 139:227 (1990)), also disclosed recently that synthetic peptides containing the HAV 35 sequence inhibited two biological processes (compaction of 8-cell-stage mouse embryos and rate of neurite

outgrowth on astrocytes) that are known to be mediated by cadherins. Effective peptides in these assays were LRAHAVDVNG and AHAVSE; PPI-containing peptides were without effect. However, Blaschuk et al. provide no
5 guidance for determining the regions flanking the HAV tripeptide that are critical for cell-cell adhesion. In the BBB disrupting peptides of the present invention detailed below, we have observed that the mere presence of the HAV sequence in a small cadherin-derived peptide
10 is not the sine qua non for a composition effective to prevent cell-cell adhesion. Indeed, it should be emphasized that neither Blaschuk et al. nor any other publication known to the present inventors suggest that cadherin sequences containing HAV or SHAVS sequences
15 would be effective in opening tight junctions and piercing blood brain barriers formed by E-cadherins in brain microvascular endothelial cells.

SUMMARY OF THE INVENTION

It has now been discovered that molecules
20 homologous to, and immunologically related to, cadherin cell adhesion molecules are present on brain and non-brain microvascular endothelial cells, such that

junctions between such endothelial cells can be reversibly opened so as to permit passage of therapeutic drugs by the use of polypeptide and antibody compositions that compete with such cell
5 adhesion molecules for binding to such cells.

It is therefore an object of this invention to provide the identity of microvascular endothelial cell adhesion molecules.

Another object of this invention is to provide DNA
10 sequences of genes, and plasmids containing same, coding for the expression of all or a cell-binding portion of microvascular endothelial cell adhesion molecules.

Yet another object of this invention is to provide
15 means to identify those sequences of cell adhesion molecules responsible for the tight binding of adjoining endothelial cells.

A further object is to provide therapeutic compositions comprising polypeptides derived from cell
20 adhesion molecules that reversibly disrupt cell-cell adhesion.

Still another object of this invention is to provide therapeutic compositions comprising polyclonal or monoclonal antibodies or fragments thereof directed
25 against endothelial cell adhesion molecules, or against polypeptides representing cell binding regions thereof, that reversibly disrupt endothelial cell-cell adhesion.

Yet another object of this invention is to provide therapeutic formulations comprising therapeutic drugs
30 conjugated with blood-brain barrier-disrupting compositions of this invention, that are capable of entering the central nervous system following disruption of the blood-brain barrier.

These and other objects of this invention will
35 become clear by reference to the following description

of the invention and to the appended claims.

DESCRIPTION OF THE DRAWINGS

Figure 1 illustrates the partial cDNA sequence for bovine endothelial cell adhesion molecule homologous to chicken N-cadherin.

Figure 2 illustrates the partial cDNA sequence for bovine endothelial cell adhesion molecule homologous to mouse P-cadherin.

Figure 3 illustrates the cDNA sequence for the MDCK cell adhesion molecule homologous to mouse E-cadherin.

Figure 4 illustrates the restriction sites in the bovine endothelial cell N- (4-1 to 4-5) and P-cadherin (4-6 to 4-8) cDNA sequences and in the MDCK E-cadherin (4-9 to 4-14) cDNA sequence.

Figure 5 shows the staining of a mouse brain thin section by an antibody raised against a fusion protein derived from amino acids 9-96 of MDCK E-cadherin containing an HAV region.

Figure 6 is a repeat of the experiment of Fig. 5, except that the antibody was raised against the entire E-cadherin molecule.

Figure 7 illustrates the effects of an 18-mer HAV-containing polypeptide on the resistance of tight junction monolayers of MDCK epithelial cells.

Figure 8 illustrates the effects of 11-mer and 18-mer HAV-containing polypeptides on the resistance of tight junction monolayers MDCK epithelial cells.

Figure 9 illustrates the effects of 11-mer and 18-mer HAV-containing polypeptides on the resistance of tight-junction monolayers of brain microvascular endothelial cells.

DETAILED DESCRIPTION OF THE INVENTION

It has now been discovered that cell adhesion molecules with characteristics of cadherins are present on the surfaces of brain capillary endothelial cells and of microvascular endothelial cells of non-brain origins. The present invention is based on the discovery that a polypeptide composition comprising cell binding domains of endothelial cell adhesion molecules may compete against such molecules for binding to such cells, such that by this means the junctions between such cells could be reversibly opened, thereby permitting penetration by therapeutic agents. The present invention also discloses that polyclonal or monoclonal antibodies (or fragments thereof) raised against endothelial cell adhesion molecules or cell-binding domains thereof may also compete for endothelial cell surface binding sites, and, by this means, reversibly disrupt junctions between endothelial cells, thereby permitting entry into the central nervous system of therapeutic agents.

In order to obtain compositions useful for disrupting tight junctions between microvascular endothelial cells, the cell adhesion molecules responsible for such junctions were identified.

The endothelial cell cadherins disclosed herein exhibit one or more of several characteristics of E-, P- and N- cadherins, including: characteristics of a transmembrane integral protein, with cytoplasmic, hydrophobic plasma membrane, and extracellular regions; intraspecies DNA sequence homologies of greater than about 50% for the entire molecule; immunological cross-reactivity with antibodies raised against non-endothelial cell cadherins; and containing cell-binding domains. "Immunologically related to" means that these cadherin-like molecules cross-react with antibodies

raised against non-endothelial cell cadherins.

E-cadherin-like molecules were localized in brain by immunofluorescence. Cryostat sections of mouse brain were labeled with a rabbit antibody prepared
5 against E-cadherin, and then with fluorescein isothiocyanate-conjugated goat anti-rabbit immunoglobulin. There is clear labeling of a capillary in brain sections as shown by immunofluorescence microscopy. Endothelial cells in liver and kidney were
10 not stained by this procedure.

cDNAs coding for the expression of bovine microvascular endothelial cell (BMEC) cadherins were cloned and sequenced as described below, and the partial sequence of N-cadherin and P-cadherin are
15 disclosed herein in Figures 1 and 2, respectively. In addition, as MDCK dog kidney epithelial cells are known to employ E-cadherin to form high resistance tight junctions, and as the present invention discloses that brain capillary endothelial cell adhesion molecules
20 include E-type cadherin, the DNA of this cadherin was also cloned; its complete DNA sequence is disclosed herein (Fig. 3).

N-, P- and E-cadherin-type clones described herein were deposited in the American Type Culture Collection
25 on September 26, 1989, and were assigned the following accession numbers:

	<u>Clone Designation</u>	<u>Accession No.</u>
	N-cadherin-type clones	
	pUC19-bNCad 10A	40667
	pUC19-bNCad 39A	40669
5	P-cadherin-type clones	
	pUC18-bPCad 3B-10	40668
	pUC19-bPCad 9B	40670
	E-cadherin-type clones	
	pBluescript MDCKECad 45-30E	40671

10 The cloning of cadherins was accomplished by taking advantage of the fact that the cadherins characterized thus far are transmembrane glycoproteins, the cytoplasmic domains of which are highly conserved, that is, are highly homologous.

15 Two degenerate oligonucleotides flanking the 42-amino acid coding region in the cytoplasmic domain were selected to serve as primers for polymerase chain reaction (PCR) using either BMEC cDNA or MDCK cDNA as templates. The PCR reactions were carried out
20 essentially according to Saiki, R. K. et al., Science, 239:487 (1988), which is incorporated herein by reference.

 The cloned PCR products from each cell type were sequenced essentially according to the method of
25 Sanger, F. et al., Proc. Nat'l. Acad. Sci. (USA), 74:5463 (1977), which is incorporated herein by reference.

 It was discovered that BMEC cadherins are of two types - one homologous to chicken N-cadherin (neuronal
30 type, see, e.g., Hatta, K., et al., J. Cell Biol., 106:873 (1988)) and the other homologous to mouse P-cadherin (placental type, see e.g., Nose, A., et al., (1987) above). It has also been found that there are two species of cadherins in MDCK cells - one homologous

to mouse E-cadherin (see, e.g., Nagafuchi, A., et al., Nature, 329:341 (1987)) and the other homologous to mouse P-cadherin (Nose, et al. (1987), above).

The PCR products were then used as probes to
5 isolate the BMEC and MDCK cadherin cDNA clones as follows. A cDNA library was constructed essentially according to Gubler et al. (Gubler, U. et al., Gene, 25:263 (1983), which is incorporated herein by reference), using poly (A)⁺RNA isolated from either
10 BMEC or MDCK cells. The cDNA was ligated via EcoRI adaptors into gt10 arms (BMEC) or ZAP^R (from Stratagene, Inc., La Jolla, CA) vector arms (MDCK). cDNA libraries containing 5×10^5 - 1.5×10^6 independent cDNA clones were screened using
15 radiolabeled PCR products (Benton, W.D. et al., Science, 196:180 (1987), which is incorporated herein by reference). Northern blot analysis (Maniatis, T. et al., "Molecular Cloning: A Laboratory Manual", Cold Spring Harbor Laboratory, Cold Spring Harbor, N.Y.,
20 1982) may be used to determine whether each cDNA species cloned hybridizes to a single mRNA species, as well as the tissue distributions of each cDNA species.

cDNA clones for each cadherin were sequenced by the method of Sanger et al. (1977) above.

25 The partial restriction maps for each cDNA clone based on their sequences are shown in Fig. 4. Some of these restriction sites were confirmed by restriction enzyme digestions, including Hind III, Pst I, Kpn I, Bgl II for N-cadherin; Pvu II, Sac I and Pst I for
30 P-cadherin; Pst I, Pvu II, BamH I, and Sac I for E-cadherin.

In order to test whether the cloned E-cadherin cDNA contains all the information necessary for cadherin function, full-length E-cadherin cDNA joined
35 to a suitable promoter may be introduced into mouse

L-cells that have very little endogenous cadherin activity (Nagafuchi, et al. (1987), supra). To test for expression of E-cadherin in transfectants derived from the introduced cDNA, transfected L-cells may be
5 tested for Ca^{2+} -dependent aggregating activity. The extent of this aggregating activity should be closely correlated with the amount of E-cadherin expressed (Takeichi, M. (1988), supra). This same technique may be used for testing cDNAs encoding bovine endothelial
10 N- and P-cadherins, according to the method of Hatta, et al. (Hatta, K., et al. (1988), supra).

In order to identify cell binding domains in, for example, MDCK E-type cadherin, L-cells may be first transfected as above with a cDNA of a size sufficient
15 to cause Ca^{2+} -mediated aggregation of transfectants. A series of deletion mutants comprising truncated cDNA species missing different regions of the extracellular domain may be prepared by restriction enzyme digestion and proper end filling or exonuclease digestion to make
20 the deletions in the proper coding frames. These deletion mutants can then be tested for their ability to express in L-cells a protein causing Ca^{2+} -dependent aggregation. By correlating a loss of aggregation with deletion of particular fragments, the regions important
25 for cell binding may be determined. A variety of polypeptides corresponding to binding regions of cadherins, as deduced from the nucleotide sequences of deleted cDNA, may be synthesized chemically using an automated peptide synthesizer such as that of Applied
30 Biosystems, Inc., Foster City, CA, or expressed by recombinant DNA methods. Effective polypeptides may be of varying lengths, depending upon the natures of junctions being disrupted and the cell adhesion molecule present.

Nucleotide, and corresponding amino acid, sequences of cadherins may be analyzed to detect homologous regions. Applying this technique to bovine endothelial cell N- and P-cadherins and to epithelial cell E-cadherin, we have determined that, in the amino acid 80 region of each of these cadherins, there is conserved a triplet HAV (His-Ala-Val) region. We have deduced that this HAV region may be a common cell adhesions recognition (CAR) sequence.

We have chemically synthesized the following polypeptides, each of which containing the HAV sequence:

	6-mer(78-83)	NH ₂ -SHAVSS-CONH ₂
	11-mer(76-86)	NH ₂ -LYSHAVSSNGN-CONH ₂
15	17-mer(74-90)	NH ₂ -YILYSHAVSSNGNAVED-CONH ₂
	18 mer(69-86)	NH ₂ -EQIAKYILYSHAVSSNGN-CONH ₂
	20-mer(71-90)	NH ₂ -IAKYILYSHAVSSNGNAVED-CONH ₂

and have tested each for efficacy in opening brain endothelial cell tight junctions in the BBB model disclosed in copending United States application Serial No. 07/413,274, and also on kidney epithelial cell tight junctions..

Polyclonal antibodies raised in rabbits and monoclonal antibodies derived from hybridomas may be generated against each of the chemically-synthesized polypeptides by standard methods. (Harlow, E., et al., "Antibodies: A Laboratory Manual", Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y., 1988; Goding, J.W., "Monoclonal Antibodies: Principles and Practice", Academic Press, N.Y. 1986). In addition, recombinant antibodies may be prepared. Fragments of antibodies, e.g., Fc, Fab, F(ab)', may be prepared by standard methods.

We have cloned and sequenced fusion proteins derived from amino acids 9-96 of MDCK E-cadherin

containing the HAV region. A polyclonal antibody prepared against this fusion protein stained rat (Fig.55) mouse brain sections as well as did an antibody raised against the entire E-cadherin (Fig. 6).

- 5 A polyclonal antibody raised against a fusion protein derived from amino acids 9-37 failed to stain brain sections. These results indicate that the key cell-binding domain of E-cadherin lies in the region of amino acids 37-96.

- 10 The ability of CAM-derived polypeptides containing cell-binding domains, and the corresponding polyclonal and monoclonal antibodies, of the invention to disrupt tight junctions may be tested in in vitro and in vivo models of high resistance tight junctions and in animal
15 models. Monolayers of MDCK dog kidney epithelial cells, that are known to contain high resistance tight junctions (Gumbiner, B., J. Cell Biol., 102:457 (1986)), can be used to test for the ability of the polypeptides and corresponding antibodies of the
20 present invention to disrupt such tight junctions.

- Polyclonal antibodies prepared as described above may also be used in conjunction with Western blotting (Old, R.W., et al., Principles of Gene Manipulation, 3d ed., Blackwell, Oxford, 1985, p. 10) and a variety of
25 tissue extracts in order to identify cell adhesion glycoproteins in such extracts.

- Another embodiment of the present invention is in drug delivery systems. Conjugates between therapeutic drugs and agents that affect cell adhesion molecule
30 function in brain capillary endothelial cells may be used to deliver therapeutic drugs to the CNS. For example, a polypeptide derived from a cell adhesion molecule that contains within its amino acid sequence a cell-binding domain, or antibodies thereto, may be
35 conjugated in biologically-active form to a therapeutic

modality. Such conjugates may have the dual effect of opening the BBB and delivering the therapeutic agent to the brain side of the BBB. Delivery of therapeutic drugs to the CNS, either alone or conjugated to agents that disrupt cell-cell adhesion, may be accomplished by administering such drugs to a subject either simultaneously with or subsequent to the administration of the agents of this invention that disrupt the tight junctions of the BBB. Examples of therapeutic modalities that may be delivered to the brain by the cell adhesion disruption compositions of this invention include Nerve Growth Factor, anti-Parkinsonian drugs, and brain enzymes known to be missing in sphingolipidoses, e.g., Tay-Sachs disease. Means of chemically conjugating protein or polypeptide carriers to therapeutic agents such that the biological integrity of the therapeutic agent is not compromised and such that the therapeutic agent is readily cleaved from the carrier by enzymes present on or within endothelial cells (e.g., amidases, esterases, disulfide-cleaving enzymes), are well known in the art. It is also apparent that these therapeutic conjugates may be delivered to endothelial cells in encapsulated form (e.g., in liposomes) or as microsuspensions stabilized by pharmacological excipients.

It is known (Jain, R.K., J. Natn'l Cancer Inst., 81:570 (1989)) that many solid tumors develop internal barriers, including high pressure zones and collapsed blood vessels, that make it difficult for blood-borne chemotherapeutic agents to reach the tumor's inner core. The barrier problem is particularly troublesome with therapeutic products drawn from the human immune system, such as monoclonal antibodies conjugated with chemotherapeutic agents, interleukin-2, interferon and activated killer T-lymphocytes, because of their large

size. Thus, in another embodiment of this invention, compositions that disrupt the junctions between endothelial cells, particularly the relatively small peptides that contain one or more cell-binding regions of cell adhesion macromolecules, may be used to enhance drug delivery to tumors with depressed blood flow.

It has been theorized that cancer cells metastasize by secreting soluble cadherins variously to open tight junctions in cells that block their movement and to prevent their being bound to such cells. We consider it likely that antibodies raised against these cadherins, which are derived from extracellular domains of the cadherins disclosed in this invention, may provide a therapeutic modality that inhibits or prevents cancer cell metastases.

In another embodiment, the compositions of this invention may also be used to provide penetration for chemotherapeutic agents of other well-known blood-tissue barriers, such as blood-testis barriers and blood-retina barriers. The latter barrier is known to prevent the efficient transport of, for example, administered antibiotics to the retina from the general circulation. The cell adhesion disrupting compositions of this invention may, thus, be used in conjunction with the administration of antibiotics to treat retinal infections.

The following examples are illustrative of several embodiments of this invention, and should not be construed in any way as limiting the invention as recited in the claims.

EXAMPLE 1

EFFECTS OF HAV-CONTAINING POLYPEPTIDES ON TIGHT JUNCTIONS OF MDCK EPITHELIAL AND BOVINE ENDOTHELIAL CELLS

The BBB model of copending U.S. Serial No. 07/413,332 was used to examine the effects of polypeptides containing the HAV region on the tight junctions of monolayers of MDCK epithelial cells and
 5 bovine capillary endothelial cells as determined by resistance measurements across the monolayers.

The polypeptide was added to the cells either from the apical side (top) or basolateral side (bottom), as shown in the following sketch.

10

APICAL

EPITHELIAL CELLS
 Gut Side

ENDOTHELIAL CELLS
 Blood Side

Blood Side

Brain Side

BASOLATERAL

15 Figure 7 illustrates the effects of various concentrations of the aforementioned 18-mer polypeptide on resistance of MDCK epithelial cells. At the lowest concentration tested, 0.5 mg/ml, resistance was markedly decreased. The polypeptide was more effective
 20 when added from the basolateral side, but at high concentrations was quite effective even when added from the apical side. These data indicate that the 18-mer is effective in making tight junctions permeable. The 20-mer was similarly effective, and a 17-mer less
 25 effective.

Figure 8 illustrates the effects of the aforementioned 11-mer and 18-mer on MDCK cell resistance when added from either the apical or
 30 of polypeptide was about 1 mg/ml. The 11-mer (as well

as the 6-mer data not shown) was virtually without effect. With the 18-mer, resistance was almost totally abolished by about 6 hours, indicating disruption of tight junctions. That the effect of the 18-mer is
5 reversible is indicated by the "wash-out" experiment. When the 18-mer was washed out of the MDCK cells at 6 hours, resistance recovered to a substantial extent over the next 21 hours. This recovery was particularly pronounced when the 18-mer had originally been added
10 from the basolateral side of the monolayers. The 20-mer produced results similar to those of the 18-mer, and the 17-mer was effective, but somewhat less so.

Figure 9 illustrates the effect of 1 mg/ml of the 11-mer and 18-mer on high resistance monolayer cultures
15 of brain endothelial cells (see copending United States Serial No. 07/413,332 for method of preparation). As with MDCK cells, the 11-mer (and the 6-mer) failed to reduce resistance values over a 48-hour period of observation. In contrast, the 18-mer (as well as the
20 20-mer) decreased resistance values markedly when added from either the basolateral or apical side, but the effect of the polypeptide was more rapid and more pronounced when it was added from the basolateral side; the 17-mer was less effective.

25 The conclusion of these experiments is that a particular set of peptides (but not all peptides) centered around the HAV region of E-cadherin are effective in opening tight junctions of brain endothelial cell blood-brain barriers, and also of
30 epithelial cells that form such junctions ("gut barrier"). Both the length and composition of the amino acid region flanking the HAV triplet thus appear to play a role in the efficacy of such compositions.

While the aforementioned embodiments represent the
35 preferred embodiments of the invention, those skilled

in the art may, without undue experimentation, devise other executions of the compositions and methods of use of this invention without departing from the concept and spirit inherent therein.

What is claimed is:

1. A composition for opening tight junctions between microvascular endothelial cells of a subject, whereby means are provided for a drug to cross the permeability barrier imposed by such junctions,
5 comprising an agent capable of reacting with at least one type of cell-bound cell adhesion molecule that would otherwise mediate tight junction formation between microvascular endothelial cells, so that cell-cell adhesion is disrupted.
2. A composition of claim 1, wherein said cell adhesion molecule exhibits at least about 50% sequence homology with a cadherin selected from the group consisting of E-cadherin, N-cadherin and P-cadherin.
3. A composition of claim 1, wherein said cell adhesion molecule is immunologically related to at least one of the group consisting of E-cadherin, N-cadherin and P-cadherin.
4. A composition of claim 1, wherein the microvascular endothelial cells are brain capillary endothelial cells.
5. A composition of claim 2, wherein said agent comprises an inhibitor of the binding to cells of said cell adhesion molecule.
6. A composition of claim 3, wherein said agent comprises an inhibitor of the binding to cells of said cell adhesion molecule.
7. A composition of claim 5, wherein said inhibitor agent comprises a fragment of said cell adhesion molecule.
8. A composition of claim 7, wherein said cell adhesion molecule fragment includes within its amino acid sequence a cell-binding domain.

9. A composition of claim 8, wherein said cell-binding domain contains an HAV amino acid sequence.


10. A composition of claim 9, wherein said amino acid sequence is



11. A composition of claim 9, wherein said amino acid sequence is



12. A composition of claim 9, wherein said amino acid sequence is



13. A composition of claim 9, wherein said amino acid sequence comprises amino acids 9-96 of E-cadherin.

14. A composition of claim 5, wherein said inhibitor agent comprises a polyclonal or monoclonal antibody directed against said cell adhesion molecule.

15. A composition of claim 5, wherein said inhibitor agent comprises a polyclonal or monoclonal antibody directed against a fragment of said cell adhesion molecule.

16. A composition of claim 15, wherein said cell adhesion molecule fragment includes within its amino acid sequence a cell-binding domain.

17. A composition of claim 16, wherein said cell-binding domain contains an HAV amino acid sequence.

18. A composition of claim 17, wherein said amino acid sequence is



19. A composition of claim 17, wherein said amino acid sequence is



20. A composition of claim 17, wherein said amino acid sequence is



21. A composition of claim 17, wherein said amino acid sequence comprises amino acids 9-96 of E-cadherin.

22. A composition of claim 5 or 6 in a pharmaceutically-acceptable vehicle.

23. A method for opening tight junctions between microvascular endothelial cells of a subject, comprising the step of administering to the subject an agent, in an effective amount and in a
5 pharmaceutically-acceptable vehicle, capable of reacting with at least one type of cell-bound cell adhesion molecule that would otherwise mediate tight junction formation between microvascular endothelial cells, so that cell-cell adhesion is disrupted and
10 whereby means are provided for a drug to cross permeability barriers imposed by such tight junctions.

24. A method of claim 23, wherein said cell adhesion molecule exhibits at least about 50% homology with a cadherin selected from the group consisting of E-cadherin, N-cadherin and P-cadherin.

26

25. A method of claim 23, wherein said cell adhesion molecule is immunologically related to at least one of the group consisting of E-cadherin, N-cadherin and P-cadherin.

26. A method of claim 23, wherein the microvascular endothelial cells are brain capillary endothelial cells.

27. A method of anyone of claims 23-25, inclusive, wherein said agent comprises an inhibitor of the binding to cells of said cell adhesion molecule.

28. A method of claim 27, wherein said inhibitor agent comprises a fragment of said cell adhesion molecule.

29. A method of claim 28, wherein said cell adhesion molecule fragment includes within its amino acid sequence a cell-binding domain.

30. A method of claim 29, wherein said cell-binding domain contains an HAV amino acid sequence.

31. A method of claim 30 wherein said amino acid sequence is



32. A method of claim 30, wherein said amino acid sequence is



33. A method of claim 30, wherein said amino acid sequence is



34. A method of claim 30, wherein said amino acid sequence comprises amino acids 9-96 of E-cadherin.


35. A method of claim 27, wherein said inhibitor agent comprises a polyclonal or monoclonal antibody directed against said cell adhesion molecule.

36. A method of claim 28, wherein said inhibitor agent comprises a polyclonal or monoclonal antibody directed against said fragment of said cell adhesion molecule.

37. A method of claim 36, wherein said cell adhesion fragment includes within its amino acid sequence a cell-binding domain.

38. A method of claim 37 wherein said cell-binding domain contains an HAV amino acid sequence.


39. A method of claim 38, wherein said amino acid sequence is



40. A method of claim 38, wherein said amino acid sequence is



41. A method of claim 38, wherein said amino acid sequence is



42. A method of claim 38, wherein said amino acid sequence comprises amino acids 9-96 of E-cadherin.

43. A drug delivery composition comprising a conjugate between a therapeutic drug and an agent capable of reacting with at least one type of a cell-bound cell adhesion molecule that would otherwise mediate tight junction formation between microvascular endothelial cells, so that cell-cell adhesion is

disrupted by said agent, whereby means are provided for said drug to cross permeability barriers imposed by such tight junctions, in a pharmaceutically-acceptable vehicle.

44. A drug delivery composition of claim 43, wherein said cell adhesion molecule exhibits at least about 50% homology with a cadherin selected from the group consisting of E-cadherin, N-cadherin and P-cadherin.

45. A drug delivery composition of claim 43, wherein said cell adhesion molecule is immunologically related to at least one of the group consisting of E-cadherin, N-cadherin and P-cadherin.

46. A drug delivery composition of claim 43, wherein the microvascular endothelial cells are brain capillary endothelial cells.


47. A drug delivery composition of any one of claims 43-45, inclusive, wherein said agent comprises an inhibitor of the binding to cells of said cell adhesion molecule.

48. A drug delivery composition of claim 47, wherein said agent comprises a fragment of said cell adhesion molecule.

49. A drug delivery composition of claim 48, wherein said cell adhesion molecule fragment includes within its amino acid sequence a cell-binding domain.

50. A drug delivery composition of claim 49, wherein said cell-binding domain contains an HAV amino acid sequence.

51. A drug delivery composition of claim 50, wherein said amino acid sequence is



52. A drug delivery composition of claim 50, wherein said amino acid sequence is



53. A drug delivery composition of claim 50, wherein said amino acid sequence is



54. A drug delivery composition of claim 50, wherein said amino acid sequence comprises amino acids 9-96 of E-cadherin.

55. A drug delivery composition of claim 43, wherein said inhibitor agent comprises a polyclonal or monoclonal antibody directed against said cell adhesion molecule.

56. A drug delivery composition of claim 43, wherein said inhibitor agent comprises a polyclonal or monoclonal antibody directed against a fragment of said cell adhesion molecule.

57. A drug delivery composition of claim 56, wherein said cell adhesion molecule fragment contains within its amino acid sequence a cell-binding domain.

58. A drug delivery composition of claim 56, wherein said cell-binding domain encompasses an HAV amino acid sequence.

59. A drug delivery composition of claim 58, wherein said amino acid sequence is



60. A drug delivery composition of claim 58, wherein said amino acid sequence is

NH₂-EQIAKYILYSHAVSSNGN-COHN₂ .

61. A drug delivery composition of claim 58, wherein said amino acid sequence is

NH₂-IAKYILYSHAVSSNGNAVED-CONH₂ .

62. A drug delivery composition of claim 58, wherein said amino acid sequence comprises amino acids 9-96 of E-cadherin.

63. A drug delivery composition of claim 43, wherein said conjugate comprises a physiologically-cleavable covalent bond.

64. A drug delivery composition of claim 43, wherein said conjugate is encapsulated within a physiologically-compatible particle.

65. A drug delivery composition of claim 64, wherein said particle comprises a liposome.

FIG. 1a.

Partial cDNA sequence for the bovine endothelial N-cadherin

GAATTGGAAC CCCTTCGTTT CATATGCAA GACTGGATTT CCTGAAGATG TGTACAGTGC	60
AGTCTTGTCC CGGGATGTGC TGGAAGGACA GCCCCTTCTC AATGTGAAGT TTAGCAACTG	120
CAATGGGAAA AGAAAAGTAC AGTATGAGAG CAGCGAGCCA GCAGATTTTA AGGTGGATGA	180
AGATGGCATG GTGTATGCCG TGAGAAGCTT CCCCCTCTCA TCTGAACACT CGAAGTTCCT	240
GATATACGCT CAAGACAAAG AGACTCAGGA AAAGTGGCAA GTAGCAGTAA AACTGAGCCT	300
CAAACCAGCC CTACCTGAGG ATTCAGTGAA GGAATCACGA GAAATAGAAG AAATAGTGTT	360
TCCAAGACAA GTGACTAAGC ACAATGGCTA CCTGCAGAGG CAGAAGAGAG ACTGGGTAT	420
CCCTCCCATC AACTTGCCAG AAAACTCCAG AGGCCCTTTT CCTCAAGAGC TCGTCAGGAT	480
CAGATCCGAT AGAGATAAAA ACCTTTCTCT GCGGTACAGC GTAACTGGGC CAGGAGCTGA	540
CCAGCCTCCA ACTGGTATCT TCATTATCAA CCCCATCTCA GGTCAAGCTGT CAGTAACCAA	600
GCCTCTGGAT CGTGAGCTGA TAGCCCGGTT TCATTGAGG GCACATGCAG TGGATATTAA	660
TGGAAACCAA GTGGAGAACC CCATCGACAT TGTCAATCAAC GTTATTGACA TGAATGATAA	720
CAGACCTGAG TTCTTACACC AGGTTTGGAA TGGGACAGTT CCTGAGGGAT CAAAGCCGGG	780
AACATATGTG ATGACGGTCA CTGCGATTGA TGCTGACGAT CCAAATGCCC TCAATGGGAT	840

1/42

FIG. 1b.

GTTGAGGTAC AGAATCCTGT CCCAGGCGCC AAGCACCCCT TCGCCCAACA TGTTTACAAT	900
CAACAATGAG ACTGGGGACA TTATCACGGT GGCAGCTGGA CTTGACAGAG AAAAAGTACA	960
ACAGTATACG TTAATAATTTC AAGCTACAGA CATGGAAGGC AATCCCACAT ATGGCCCTTTC	1020
CAACACAGCC ACGGCTGTCA TCACGGTGAC AGATGTCAAC GACAATCCTC CGGAGTTTAC	1080
TGCCATGACG TTCATGTGTG AAGTCCCTGA AAACAGGGTA GATGTCATCG TCGCTAATCT	1140
AACAGTGACA GATAAGGATC AGCCCCACAC ACCGGCCTGG AACGCCATCT ACAGAATCAG	1200
CGGTGGAGAC CCCGCCGGCC GCTTTGCCAT TCAAACTGAC CCCAACAGCA ACGACGGTTT	1260
AGTCACCGTA GTAAAACCAA TCGACTTTGA AACAAATAGG ATGTATGTCC TTTACTGTGCG	1320
TGCAGAAAAT CAAGTGCCAT TAGCCAAGGG TATTCAGCAT CCACCTCAGT CAACTGCCGAC	1380
TGTGTCTGTC ACAGTTATCG ATGTGAATGA AAATCCTTAT TTTGCCCCCA ATCCAAAGAT	1440
CATTGCGCAA GAAGAAGGCC TTCACGCCGG TACCGTGTTA ACAACGTTTA CTGCTCAGGA	1500
CCCAGATCGA TATATGCAGC AAAATATCAG ATACACCAA TTATCCGATC CTGCAAACTG	1560
GCTAAAAATA GACTCTGTGA ATGGGCAGAT AACTACCATT GCTGTTTGG ACAGAGAATC	1620
ACCGAATGTG AAAGCCAATA TATACAATGC TACTTTCCTT GCTTCTGACA ATGGAATCCC	1680
TCCTATGAGT GGAACGGGAA CACTGCAGAT CTATTTACTT GATATTAAATG ACAATGCCCC	1740

FIG. 1c.

TCAAGTGTTA	CCTCAAGAGG	CAGAGATTGG	TGAAACTCCG	GACCCCAATT	CAATTAAACAT	1800
CACAGCACTT	GATTATGACA	TTGATCCAAA	TGCTGGACCA	TTTGCTTTTG	ATCTTCCTTT	1860
GTCTCCAGTG	ACTATTAAAG	GAAATTGGAC	CATCACTCGG	CTTAATGGTG	ATTTTGCTCA	1920
GCTTAACTTA	AAGATAAAAT	TTCTTGAGGC	CGGATCTAC	GAAGTTCCAA	TCATAATCAC	1980
AGATTGGGT	AATCCTCCCA	AATCGAATAT	CTCCATCCTT	CGGGTGAAGG	TTTGCCAGTG	2040
TGATTCCAAC	GGGACTGCA	CAGATGTGGA	TCGAATTGTG	GGAGCAGGGC	TGGGCACCGG	2100
CGCCATCATC	GCCATCCTGC	TTTGCAATCAT	CATCCTGCTC	ATTCTCGTTC	TGATGTTCGT	2160
GGTATGGATG	AAACGCCGGG	ATAAAGAACG	CCAGGCCAAA	CAACTTTTAA	TTGATCCAGA	2220
AGATGATGTA	AGAGATAATA	TTTTAAATA	TGATGAAGAA	GGTGGAGGAG	AAGAAGACCA	2280
GGACTACGAT	TTGAGCCAGC	TCCAGCAGCC	TGATACGGTA	GAGCCAGATG	CCATCAAGCC	2340
AGTTGGAATC	CGACGGTTGG	ATGAGAGGCC	CATCCATGCG	GAGCCCCAGT	ACCCGGTTCCG	2400
ATCTGCAGCC	CCACACCCAG	GGACATCGG	GGACTTCATT	AATGAGGGCC	TTAAAGCTGC	2460
TGACAAACGAT	CCCACCGCTC	CGCCCTACGA	CTCCCTCTTA	GTCTTTGACT	ATGAAGGCAG	2520
TGGCTCCACG	GCCGGGTCTT	TGAGCTCCCT	TAATTCCTCC	AGTAGTGGAG	GTGAGCAGGA	2580
CTATGACTAT	CTGAACGACT	GGGGGCCCCG	CTTCAAGAAA	CTCGCTGACA	TGTACGGTGG	2640

SUBSTITUTE SHEET

FIG. 1d.

AGGTGATGAC TGAAGCTTCAG GGTGAACTTG GTTTTTGGAC AAGTACAAAC AATTGCAACT 2700
GATATTCCCA AAAAGCATTC AGAAGCTAGG CTTTAACTTT GTAGTCTACT AGCACAGTGC 2760
TTGCTGGAGG CTTTGGCAGA GGCTGCAAAC CAATTGGGC TCAGAGGGA TATCGGTGAT 2820
CCAATACTGT TTGGAAAACA CTGAGCTCAG TTACACTTGA ATTTTACAGT ACAGAAGCAC 2880
TGGGATTTTA TGTGCCCTTTT TGTACCTTTT TCAGATTGGA ATTAGTTTTA TGTTTAAGGC 2940
TTTAAATGGTA CTGATTCTG AATGATAAG TAAAAGACAA AATATTTTGT GGTGGGAGCA 3000
GTAAGTTAAA CCATGATATG CTTCGACACG CTTTGTGTAC ATCGCATTTG CTTTATTATA 3060
AAATATGGAA TTAAACAGAC AAACCAACCA CTCATGGAGC AATTTTATTA CCTTGGGGGC 3120
TGAGACCATG AGATTGAAA ATGTACATTA TTTCTAGTTT TAGACTTTAG TTTCTTGTTT 3180
TGTTTTTTTT TTCCACTAAA ATCTTAAAC TTACGCAGCT GGTGCAAAAT AAAGGGAGTT 3240
TTCATATCAC CAATTGTAG CAAAATTGAA TTTTTCATA AACTAGAAATG TTAGACACAT 3300
TTTGGTCTTA ATCCATGTAC ACTTTTTTAT TTACTGTATT TTTTCCACTT CACTGTAAA 3360
ATGGTATGTG TACATAATGT TTTATGGCA TAGTCTATGG AGAAGTGCAG AAACCTCAGA 3420
ACATGTGTAT GTATTATTG GACTATGGAT TCAGGTTTTT TGCATGTTTA TATCTTTTCTG 3480
TATGGATAAA GTATTTACAA AACAAAGTGA CATTGATTC AATTGTTGAG CTGTAGTTAG 3540
AATACTCAAT TTTTAAATTTT TTAATTTTTT TTAATTTTTA TTTTCTCTTT TTGTTTGGG 3600

SUBSTITUTE SHEET

AGGGAGAAAA GTTCTTAGCA CAAATGTTTT ACATAATTG TACCAGAAAA AACAAAAAA 3660
AAAGGAAAGA CAAGAAATGA AAGGGGTGAC CTGACACTGG TGTACTACT GCAGTGTGTG 3720
TTTTTAAAAA AAAATGAAAA AAAAAAGCT TTTAAACTGG AGAGACTTCT GACAAACAGCT 3780
TTGCCCTCTGT ATTGTGTACC AGAATATAAA TGATACACCT CTGACCCCCAG CGTCTCTGAAT 3840
AAAATGCTAA TTTTGAAAAA AAAAAAAA AAAA 3875

FIG. 1e.

FIG. 2a.

partial cDNA sequence for the bovine endothelial P-cadherin

GAATTGGAAC CCCTTCGCTG AGAACACAGT GAGCCACGAG GTGCAGAGGC TGACAGTGAC 60
TGATCTGGAC GCCCCTAACT CACCAGCATG GCGTGCCACC TACCGCATCG TGGGAGGTGA 120
CAACGGGGAC CATTTTACCA TCACTACTGA CCCCAGAGC AACGAGGTA TCCTGACCAC 180
CCAGAGGGC TTGGATTTTG AGGCCAAAAC CCAGCACACC CTGTACGTCG AAGTGATCAA 240
CGAGGTTCCT TTTGTGGTGA AACTCCCCGAC CTCCACAGCC ACCGTAGTGG TCCTCGTGGA 300
GGATGTGAAT GAGCCACCCG TGTGTGTCCC CCCGTCCAAA GTCATCGAAA TCCAGGAGGG 360
CATCTCCACT GGGGAGCCTA TTTGTGCCTA CACTGCACGG GACCCAGACA AGGGGAGTCA 420

SUBSTITUTE SHEET

FIG. 2b.

6/42

GAAGATCAGT TACCACATCC TGAGAGACCC AGCAGGGTGG CTAGCGATGG ACCCAGACAG 480
TGGACAAGTC ACTGCCGCAG GGGTCTTGGA CCGTGAGGAT GAGCAGTTTG TGAGAAACAA 540
CATCTACGAA GTCATGGTCT TGGCCACAGA TGATGGGAGC CCTCCCACCA CTGGCACAGG 600
GACCCCTCCTG CTAACACTGA TGGACATCAA TGACCACGGT CCGGTCCCCG AGCCCCGTCA 660
GATCACCATC TGCAACCAA GCCCTGTGCC CCAGGTGCTA AACATCACAG ACAAGGACTT 720
GTCCCCCCAC ACTGCCCTT TCCAGGCCA ACTCACACAT GACTCGGACG TCTATTGGAC 780
AGCAGAAATC AACGAGAAAG GAGACGCAGT AGCCTTGTC CTGAAGAAAGT TCCTAAAGCA 840
AGCGGAATAC GATGTGCACC TTTCCCTGTC CGACCACGGC AACAAGGAAC AGCTGACAGT 900
GATCAGAGCC ACCGTGTGTG ACTGCCACGG CAACATGGTG ACCTGCCGGG ACCCTGGAC 960
GTGGGGTTTC CTCCTCCCCA TCCTGGGTGC TGCCCTGGCT CTGCTGCTCC TTCTGCTGGT 1020
GCTCCTATTC TTGGTGAGAA AGAAACGGAA GATCAAGGA CCCCTTCTCC TCCCAGAAGA 1080
TGATACCCGT GACAACGTCT TCTACTACGG CGAAGAGGGG GGTGGCGAGG AGGACCAGGA 1140
CTATGACATC ACCCAGCTCC ACCGGGGTCT GGAGGCCCGG CCTGAGGTGG TTCTCCGCAA 1200
CGATGTGGCA CCATCCTTCA TCCCCACACC CATGTACCGT CCTCGGCCAG CCAACCCAGA 1260
TGAAATCGG AACTTCATCA TTGAGAACCT GAAGGCAGCC AACACAGACC CCACGGCCCC 1320

SUBSTITUTE SHEET

GCCTACGAC TCCCTGTTGG TGTTGACTA TGAGGGCAGT GGCTCCGATG CCGCCTCTCT 1380
GAGCTCGCTC ACCTCCTCAA CCTCTGACCA GGACCAAGAC TACAACTATC TGAATGAGTG 1440
GGCAGCCGC TTCAAGAAGC TGGCGGACAT GTACGGCGGG GGCAGGACG ACTAGGACTC 1500
CCTAAACGCC GGGCTGCAGC AGCGTCTCCA AGGGGTCACT ATCCCCACGT TGGCCAAGGA 1560
CTTTGCAGCT TGTGAGAAT TGGCCTTAGC AACTTGGAGG GAAGAGGCCT CGAAACTGAC 1620
CTCAAAGGGG CAGGTCTCTA TGCCTTTCAG AACGGAGGAA CGTGGGCAGT TTGATTTCAA 1680
CAGTGAGCAC CTCTTAGCCT AAGCCAGGGC TGCTCAATT CTGGGAGTCT CCTCGCTACC 1740
ATAAAATGCT CAGCGCTGGG TCCTGGGTTT TGA CTGACTC TGA CTTTCCC ATGATGGCTT 1800
TTGCTCTGGA ATGGACCCTT CTCCTTAGTA ACAGGCCCTCT TACCACAATC TTCGTTTTTT 1860
TTTTTTTAAAT GCTGTTTTCA AAAAGTGAGA GGCAGGTCCT CAACCACCCC CTGGAGCGCT 1920
CCAGAAGCCC AGCGTGCCC TCATGCATTT CTCTGTGTC TCTTGGCCCC CAGACCTCCT 1980
GTTTGATGG ATA ACTGCAT TTTTATACTG AGCAGTCTA AGTGGTCCTT TATTTTTTAT 2040
TTTCCCTATC GAGTGCTGTA GATGAAGAGT GATGACAATC CTGTAAATGT ACTAGAACTT 2100
TTTTATATAA GGA ACTTTTT CCCAAAAAAA AAAAAAAA AAAAAC 2156

FIG. 2c.

SUBSTITUTE SHEET

FIG. 3a.

cDNA sequence for MDCK E-cadherin

CGGGCACCTG	TGATTGCGG	AAGTCCTGCC	GCCTCGCGCC	GCCTCGCGCC	CGGCTCTCGA	60
CCCCCGCCCC	CCATGGGCCC	TCGGTACGGC	GGCGCCCCCG	CGCTCCTGCT	CCCGCTGCTG	120
CTGCTGCTGC	AGGTCTCATC	GGGCTCTGTC	CAAGAGCCGG	AGCCCTGCCG	CCCTGGCTTT	180
GGCGCTGACA	GCTACACGTT	CACCGTGCCC	CGCGACACT	TGGAGAGAGG	CCGTGTCCTG	240
GGCAGGGTGA	GTTTGAAGG	ATGCACCGGT	CTACCTAGGA	CAGCCTATGT	TTCTGATGAC	300
ACCCGATTCA	AAGTGGGCAC	AGATGGTGTG	ATTACAGTCA	AGCGCCTCT	ACAACTTCAT	360
AAACCAGAGA	TAAGTTTCT	TGTCCATGCC	TGGGACTCCA	GCCGCAGGAA	GCTCTCCACC	420
AGAGTTAGGC	TGAAGGCAGC	GACGCACCAC	CACCACCACC	ATCATGATGC	TCCCTCTAAA	480
ACCCAGACAG	AGGTGCTCAC	ATTTCCTCAGT	TCCCAGCATG	GACTCAGAAG	ACAGAAGAGA	540
GACTGGGTTA	TCCCTCCTAT	CAGCTGCCCG	GAAAACGAGA	AAGGCCCATTT	TCCTAAAAAC	600
CTGGTTCAGA	TCAAGTCTAA	CAGGGACAAA	GAAATCAAGG	TTTTCTACAG	CATCACTGGC	660
CAAGGAGCTG	ACGCACCTCC	TGTTGGTGTG	TTTATTATTG	AAAGAGAAAC	AGGATGGCTG	720
AAGGTGACTG	AGCCTCTGGA	TAGAGAACAA	ATTGCTAAGT	ACATTCTCTA	CTCTCATGCC	780
GTATCTTCTA	ATGGGAATGC	GGTTGAAGAC	CCAATGGAGA	TCGTGATCAC	GGTGACAGAT	840

FIG. 3b.

CAGAATGACA ACAAGCCCGA GTTCACCCAG GCAGTCTTCC AAGGATCTGT CACGGAAGGT 900
GCCCTTCCAG GCACCTCTGT GATGCAGGTG ACAGCCACAG ATGCGGATGA TGATGTGAAT 960
ACCTACAACG CTGCCATCGC TTACAGCATC CTCACACAAG ACCCCCTCCT GCCTAGCAGC 1020
ATGATGTTCA CTATCAACAA GGACACAGGA GTCATCAGCG TGCTCACCAC TGGGCTGGAC 1080
CGAGAGGGTG TCCCCATGTA CACCTTGGTG GTTCAGGCTG CTGACCCTGCA AGCGGAAGGC 1140
TTAACTACAA CTGCAACAGC TGTGATCACA GTCACCTGACA TCAATGATAA CCCCCCATC 1200
TTCAACCCAA CCACGTACCA GGGACGGGTG CCTGAGAAAC AGGCTAAAGT CGAAATCGCT 1260
GTA CTCAAAG TGACGGATGC TGATGTCCCC GATACCCCGG CCTGGAGGGC TGTGTACACC 1320
ATATTGAACA ATAACAATGA TCAATTGTGT GTCACCACAG ACCAGTAAC TAACGACGGC 1380
ATTTTGAAAA CAACTAAGGG CTGGGATTTT GAGGACAAGC AGCAGTATGT CTTGTACGTG 1440
ACTGTGGTGA ACGTGACCCC GTTTGAGGTC ATCCTCTCCA CCTCCACAGC CACTGTCACT 1500
GTGGACGTGG AAGATGTGAA TGAAGCCCCC ATCTTCATCC CTGCCCCAAA GGAGTGTCA 1560
ATCCCTGAAG ACTTTGGTGT GGGCCAGGAA ATCACAATCCT ACACCGCCGA GGATCCAGAT 1620
ACATATATGG AACAGAGGAT AACGTATCGG ATTTGGAGGG ATGCTGCCGG TTGGCTGGAG 1680
GTTAATCCAG AATCTGGTGC CATTTTCACT CGGGCTGAGC TGGACAGAGA GGATTTTGAG 1740

9/42

SUBSTITUTE SHEET

FIG. 3c.

10/42

CACGTGAAGA ATAGCACGTA TGAAGCCCTC ATTATAGCCA TTGACTTCGG TTCTCCAGTT 1800
GCTACTGGAA CGGGAACCTCT TCTACTGGTC CTCTCTGATG TGAATGACAA TGGCCCCATT 1860
CCAGAACCTC GAAATATGGA CTTCTGCCAG AAAAACCAC AGCCTCATGT CATCAACATC 1920
ATTGATCCAG ATCTTCCCCC CAACACATCT CCCTTCACAG CAGAACTAAC ACACGGCGCA 1980
AGTGTCAACT GGACCATCGA GTACAATGAC CCAGCTCGTG AATCTCTAAT TTTGAAGCCA 2040
AAGAAAACCT TAGAGTTGGG TGACTACAAA ATAAATCTCA AGCTCACAGA TAACCAGAAC 2100
AAGGACCAGG TGACCACCCT ATATGTGTTT GTGTGGGACT GCGAAGGTGT CGTCAACAGC 2160
TGCAAGAGGA CGGCGCCTTA CGCCGAAGCA GGCTTGCAGG TTCCTGCCAT CTGCGGCATT 2220
CTCGGAGGAA TCCTCGCTCT ACTAATCCTG ATTCTGCTGC TTCTGCTATT TGTTCGGAGG 2280
AGAAGGGTGG TCAAAGAGCC CTTACTTCCC CCAGAAGATG ACACCCGGGA CAATGTTTAT 2340
TACTATGATG AAGAAGGAGG TGGAGAGGAG GATCAGGACT TTGACTTGAG CCAGTTGCAC 2400
AGGGGCCCTG ATGCTCGGCC TGAAGTGACT CGCAATGATG TGGCCCCAAC CCTCCTGAGT 2460
GTGCCCCAGT ATCGGCCCCG CCCTGCCCAAT CCTGATGAAA TTGGAAACTT TATTGATGAA 2520
AACCTGAAGG CAGCGGACAC TGACCCCTACT GCTCCTCCTT ATGACTCTCT GCTCGTGT 2580
GACTATGAAG GAAGCGGTTT TGAAGCTGCT AGTCTGAGCT CCTTGAACCT CTCAGAGTCA 2640
GACCAAGACC AGGACTATGA CTACCTGAAT GAATGGGGCA ATCGCTTCAA GAAGCTGGCG 2700

SUBSTITUTE SHEET

FIG. 3d.

GACATGTATG	GAGGTGGCGA	GGACGACTAG	GGGACTTGAG	ACAAATGAAG	ATGAGTCCTT	2760
ATACCATGTG	GTAGAAAATG	CGGAGGTGAC	TGTTTTCAGC	TCCCTTCATC	TGAGAGGAAT	2820
TTCTGGAGAA	GAGAAAATGC	ACAGTGATAT	ATAGTTAGGA	TAGTTAGGAT	TTCTACTTTA	2880
TAGATCTAAT	CTGTGTGTTT	GTTAGAACGA	TTTTGACCCTA	TTCTTTGAAG	CTTTTTTTTC	2940
TTTCTTTCAT	CATTCTTTAA	ATGGTGATGC	TGTCCAAAAG	ACCCCCACA	TGTTTATATT	3000
TCAAAAGAAT	AGCTAAAGCC	TCCAGAAGGT	TCTGCTAGCA	ATTTTCGAGAT	TGCCTTATTG	3060
ACTTGCTCTCA	TTTTTTTAA	GGAAGGTAGG	GCTAAACTAC	CCTATTGTGT	TTGTGTGTGT	3120
GTGTGTGTAT	GTGTAATTAT	TTTTAATTTG	TGTTCTTTTT	TCTCCTATCA	CTGCAC TGGT	3180
GTCCCGTGTT	CTAATAACCA	CTCTTAACTC	CTTCTGAACT	TACATTGCCT	CAGACAGGAG	3240
TTCTCTGCTG	CAGAAATTAT	TGGGCCCTTT	CAGGATAAGA	GACTTGGTCT	TAGTTTGATG	3300
GTAGTGTGAC	TGGGTATTAT	GGACTCGTAA	GGACTTTAGT	GGTCTCCTT	TTTTTATTCC	3360
TAAGTACATA	AATTGAAATT	CATATCCATC	CACTGACTTG	TTCTGCATTA	AGTGTGTTTG	3420
TCATGTGAC	GTCATTATTG	GGCTACTTTG	GTTCGAAACA	AGGAGCATTG	ACCAGAAAAG	3480
GTGGTGAATT	TTTCAGGTGCC	ACTCAACTTC	TAATGTTTAC	TTATCACTCA	AACAGAAGAG	3540
TGATCTATTC	TGACGTTTAG	CGTAGTGCCT	GCAGTGCTGC	AGCCAAAAGAT	TGAAGGCCGA	3600

11/42

12/42

TTGTCAAAGC CAAGGCAAC ATGAAAAATG GACTTGGAGG TGGCAGGCGG GATGGGTCAT 3660
TGAGCCCTGGC GTTTTAGCAA ACTGATGCTG AGGATAACTG AGGTGGCTCT ACCTCTAGTC 3720
CTGAAAATTC TGAAGAAATG AAGAAATCCG ACAAGTGTGT CCTATCGCGA TCCTTAGGTC 3780
ACAGTTTGTA CCTGAGGCCA AGAATCCCCA GGTGCCCTGCT TTTGTTAATG TCTACCGAAA 3840
ATGCAGCCTG ATCTGGACTC AGGTGCCCCA ATTCTAAGTG TGCATAGAAA ACTGACAATA 3900
TTAGGAAATT CTTTTTCCCC CCTTAGGAGC AGGAAGAAAA TATGACCCCTA AAGGGTTTTC 3960
GCAAGGGAA GGTGGGGAGA GCTTTGACTT GGATTTTTTT TAAATTGAAA TGTGAACTTC 4020
AAGGAACTTT TGACAACCAT GGGAAATAAT TTTATCTTAA ATTGCTTTAC TGCTGTCTCAG 4080
CTGTTTTTCA AAGAAAAAAA AAATCATCCC TGCAATCACT TCTTGGAAAT GTCTTGATTT 4140
TTCAGCAAAT TAAACTCTAA TTAGTCCTG TATAGAGAAT GTTAATGTAG TTTTGAGTGT 4200
ATATGTGTGT GGTACGGAT AATTTTGAT TTTCTTTAGG TCTGAAAAAG GAAAAACAAT 4260
TAAGCTGCGA AAATTCTTAA ATATTCATTT TTATAAATTT TATTAAAGAA TTTTGTGTTAAA 4320
AAAAAAAAAA AAA 4333

FIG. 3e.

SUBSTITUTE SHEET

N-cadherin restriction map

[illegible]

FIG. 4b.

BspMI
 PstI
 TCCAGACAAGTGACTAAGCACAAATGGCTACCTGCAGAGGCCAGAAAGAGAGACTGGGTTAT 420
 SstI
 SacI
 HgiAI
 Bsp1286
 BanII
 EcoO109
 EaeI
 DraII
 CCTCCCATCAACTTGCCAGAAAACTCCAGAGGGCCCTTTTCTCAAGAGCTCGTCAGGAT 480
 XhoII
 AlwNI
 CAGATCCGATAGAGATAAAACCTTTCTCTGCGGTACAGCGTAACTGGGCCAGGAGCTGA 540
 PvuII
 CCAGCCTCCAACCTGGTATCTTCATTATCAACCCCATCTCAGGTCAGCTGTCAAGTAACCAA 600
 BstXI
 NspHI
 Bsp1286
 AseI

FIG. 4c.

15/42

GCCTCTGGATCGTGAGCTGATAGCCCGGTTTCATTTGAGGGCACATGCAGTGGATATTAA 660

Tth111I

TGGAAACCAAGTGAGAAACCCCATCGACATTGTCATCAACGTTATTGACATGAATGATAA 720

SauI

Eco81I

Bsu36I

AlwNI

CAGACCTGAGTTCTTACACCAGGTTTGAATGGGACAGTTCCTGAGGGATCAAAGCCGGG 780

NdeI

AACATATGTGATGACGGTCACTGCGATTGATGCTGACGATCCAAATGCCCTCAATGGGAT 840

HaeII

BbeI

NarI

BanI

EcoNI AhaII

NspHI

AflIII

GTTGAGGTACAGAAATCCTGTCCCAGGCGCCCAAGCACCCCTTCGCCCAACATGTTTACAAT 900

PvuII

CAACAATGAGACTGGGACATTATCACGGTGGCAGCTGGACTTGACAGAGAAAAAGTACA 960

FIG. 4d.

16/42

AccI		NdeI	
ACAGTATACGTTAATAATTCAAGCTACAGACATGGAAGGCAATCCCACATATGGCCTTTC			1020
		BspMII	
	HincII	AccIII	
CAACACAGCCACGGCTGTCATCACGGTGACAGATGTCAACGACAATCCTCCGGAGTTAC			1080
TGCCATGACGTTCTATGGTGAAGTCCCTGAAACAGGGTAGATGTCAATCGTCGCTAATCT			1140
	Cfr10I		
AACAGTGACAGATAAGGATCAGCCCCACACACCGGCGCTGGAAACGCCCATCTACAGAATCAG			1200
NaeI			
Eco52I			
EagI			
Cfr10I			
CGGTGGAGACCCCGCGCGCTTTGCCATTCAAACTGACCCCAACAGCAACGCGGTTT			1260
AGTCACCGTAGTAAACCAATCGACTTTGAAACAAATAGGATGTATGTCCTTACTGTCGC			1320
PstI	StyI	HincII	
TGCAGAAATCAAGTGCCATTAGCCAAGGGTATTCAGCATCCACCTCAGTCAACTGCGAC			1380

SUBSTITUTE SHEET

17/42

[illegible]

SUBSTITUTE SHEET

FIG. 4f.

18/42

PflMI		
CACAGCACTTGATTATGACATTGATCCAAATGCTGGACCATTTGCTTTTGATCTTCCTTT		1860
CelII		
GTCTCCAGTGA CTATTAAGAGAAATTGGACCATCACTCGGCTTAATGGTGATTGTGCTCA		1920
XhoII		
GCTTAACCTTAAAGATAAAATTCTTGAGGCCGGGATCTACGAAGTTCCAATCATATCAC		1980
AGATTCCGGGTAATCCTCCCAAATCGAATATCTCCATCCTTCGGGTGAAGGTTTGCCAGTG		2040
Cfr10I		
Bsp1286		
BanI		
BanI		
TGATTCCAACGGGGACTGCACAGATGTGGATCGAATTGTGGGAGCAGGGCTGGGCACCGG		2100
HaeII		
BbeI		
NarI		
AhaII		

SUBSTITUTE SHEET

19/42

FIG. 4g.

CGCCATCATCGCCATCCTGCTTTGTCATCATCTCTGCTCATTTCTCGTTCTGATGTTTCGT 2160
GGTATGGATGAAACGCCCGGGATAAAGAACGCCAGGCCAAACAACTTTTAAATTGATCCAGA 2220
AGATGATGTAAGAGATAATATTTTAAATATGATGAAGAAGGTGGAGGAGAAGACCA 2280
GGACTACGATTTGAGCCAGCTCCAGCAGCCTGTATACGGTAGAGCCAGATGCCATCAAGCC 2340
AGTTGGAATCCGACGGTTGGATGAGAGGCCCATCCATGCGGAGCCCCAGTACCCGGTTCCG 2400
ATCTGCAGCCCCACACCCAGGGGACATCGGGGACTTCATTAAATGAGGGCCCTTAAAGCTGC 2460
TGACAAAGATCCCACCGCTCCGCCCTACGACTCCCTCTTAGTCTTTGACTATGAAGGCAG 2520
TGGCTCCACGGCGGCTCCTTGAGCTCCCTTAAATTCCTCCAGTAGTGAGGTGAGCAGGA 2580

DraI
SspI AhaIII
EaeI
Bsp1286
BanII
PstI
EcoO109
EaeI
AseI
DraII
SstI
SacI
HgiAI
Eco52I
EagI
DraII
Bsp1286
BanII

SUBSTITUTE SHEET

FIG. 4h.

20/42

Bsp1286
 BanII
 ApaI
 Eco0109
 DraII
 Eco0109
 EaeI
 DraII
 NspHI
 AflIII
 CTATGACTATCTGAACGACTGGGGGCCCCCGCTTCAAGAACTCGCTGACATGTACGGTGG 2640
 AGTGATGACTGAACCTTCAGGGTGAACTTGGTTTGGACAAGTACAAACAATTGCAACT 2700
 BsmI
 ACCI
 GATATCCCAAAAGCATTTCAGAAGCTAGGCTTTAACTTTGTAGTCTACTAGCACAGTGC 2760
 AlwNI
 Bsp1286
 BanII
 TTAGCTGGAGGCTTTGGCAGAGGCTGCAAAACCAATTGGGCTCAGAGGGAATATCGGTGAT 2820
 SstI
 SacI
 HgiAI
 Bsp1286

SUBSTITUTE SHEET

21/42

FIG. 4i.

BanII
|
CCAA TACTGTTTGGAAAACACTGAGCTCAGTTACACTTGAATTTTACGTACAGAACGAC 2880
TGGGATTTTATGTGCCCTTTTGTACCTTTTTCAGATTGGAATTAGTTTATGTTTAAAGGC 2940

SspI
|
TTTAATGGTACTGATTTCTGAAATGATAAGTAAAGACAAAATATTTTGTGTGGAGCA 3000
GTAAGTTAAACCATGATATGCTTCGACACGCTTTTGTACATCGCAATTTGCTTTTATTAA 3060

StyI
|
AAATATGGAAATTAAACAGACAAACCAACCTCATGGAGCAATTTTATTACCTTGGGGC 3120
TGAGACCATGAGATTGGAAAATGTACATTATTTCTAGTTTTAGACTTTAGTTTCTTGTTT 3180

BstXI
|
TGTTT TTTTCCACTAAATCTTAAACTTACGAGCTGGTTGCAATAAAGGAGTT 3240

PvuII
|
TTCATATCACC AATTGTAGCAAAATTTGAATTTTTCATAAACTAG AATGTAGACACAT 3300
TTTGGTCTTAAATCCATGTACACTTTT TTTTATTACTGTATTTTTCCTTCACTGTA AAA 3360
ATGGTATGTACATAATGTTTATTGGCATAGTCTATGGAGAA GTGCAGAAACTTCAGA 3420

XmnI
|

SUBSTITUTE SHEET

FIG. 4j.


22/42

NspHI			
AflIII	NspHI		
ACATGTGATGTATTATTGGACTATGGATTCAGGTTTTTTGCAATGTTTATATCTTTCGT			3480
TATGGATAAAGTATTTACAAAACAAAGTGACATTGTGATTCAATTGTTGAGCTGTAGTTAG			3540
AATACTCAATTTTAAATTTTAAATTTTAAATTTTAAATTTTAAATTTTCTCTTTTGTGTTGGGG			3600
AGGAGAAAAGTCTTAGCACAAATGTTTACATAATTGTGTACCAAAAAACAAAAA			3660
	BstEII	PstI	
AAAGGAAAGACAAAGAAATGAAAGGGGTGACCTGACACTGGTGGTACTACTGCAGTGTGTG			3720
	DraI	DraI	
	AhaIII	AhaIII	
		HindIII	
TTTTTAAAAAAATGAAAAAAAGCTTTTAAACTGGAGAGACTTCTGACAAACAGCT			3780
TTGCCTCTGTATTGTGTACCAGAATATAAATGATACACCTCTGACCCCGTCTCTGAAT			3840
AAAATGCTAATTTTGGAAAAAATAAAAAA			3875

SUBSTITUTE SHEET

FIG. 4k.

P-cadherin restriction map



23/42

SUBSTITUTE SHEET

FIG. 4I.

24/42

NheI
 BstXI
 PflMI
 480
 GAAGATCAGTTACCATCCTGAGAGACCCAGCAGGGTGGCTAGCGATGGACCCAGACAG
 TGGACAAAGTCACTGCCCGCAGGGGTCTTGGACCGTGAGGATGAGCAGTTTGTGAGAAACAA 540
 Bsp1286
 BalI
 BanII
 600
 CATCTACGAAGTCATGGTCTTGGCCACAGATGATGGAGCCCTCCCACTGGCACAGG
 Eco0109
 DraII
 Bsp1286
 BanII
 660
 GACCTCCTGCTAACACTGATGGACATCAATGACCACGGTCCGGTCCCGAGCCCGTCA
 Bsp1286
 720
 GATCACCATCTGCAACCAAGCCCTGTGCCCCCAGGTGCTAAACATCACAGACAAGGACTT
 EaeI
 AatII
 AhaII
 780
 GTCCCCCAGACTGCCCTTTCCAGGCCCAACTCACACATGACTCGGACGTCTATTGGAC
 HincII
 XmnI
 840
 AGCAGAAGTCAACGAGAAAGGAGACGCAGTAGCCCTGTCCCTGAAGAAAGTTCCCTAAAGCA

SUBSTITUTE SHEET

FIG. 4m.

25/42

HgiAI Bsp1286 ApaI		PvuII	900
AGCGGAATACGATGTGCACCTTCCCTGTCCGACCACGGCAACAAGGAACAGCTGACAGT			
BclI	DraIII	BspMI EcoO109 DraII	960
GATCAGAGCCACCGTGTGACTGCCACGGCAACATGGTGACCTGCCGGGACCCCTGGAC			
GTGGGGTTTCCTCCTCCCATCCTGGGTGCTGCCCTGGCTGCTGCTCCTTCTGCTGCTGT			1020
HgiAI Bsp1286		XmnI	1080
GCTCCTATTCTTGGTGAGAAAGAAACGGAAGATCAAGGAACCCCTTCTCCTCCCAAGA			
Tth111I			1140
TGATACCCGTGACAAACGTCTTCTACTACGGCGAAGAGGGGGTGGCGAGGAGGACCAGGA			
CTATGACATCACCAGCTCCACCGGGTCTGGAGGCGCCCGGCTGAGGTGGTTCTCCGCA		SmaI Eco81I Bsu36I	1200

SUBSTITUTE SHEET

FIG. 4n.

26/42

BanI
|
CGATGTGGCACCATCCTTCATCCCCACACCCATGTACCGTCCTCGGCCAGCCAAACCAGGA 1260
TGAAATCGGCAACTTCATCATTTGAGAACCTGAAGGCAGCCCAACACAGACCCCAACGGCCCC 1320
GCCCTACGACTCCCTGTGTGTTTCGACTATGAGGGCAGTGGCTCCGATGCCGCCTCTCT 1380

SstI
SacI
HgiAI
Bsp1286
BanII
|

GAGCTCGCTCACCTCCTCAACCTCTGACCAGGACCAAGACTACAATCTCTGAATGAGTG 1440

NspHI
AflIII
| |

GGGCAGCCGCTTCAAGAAGCTGGCGGACATGTACGGGGGGCCAGGACGACTAGGACTC 1500

PstI StyI
| |
CCTAAACGCCGGCTGCAGCAGCGTCTCCAAGGGTCACTATCCCCACGTTGGCCAAGGA 1560

StuI
EaeI
| |
CTTTGCAGCTTGTGAGAAATTGGCCCTTAGCAACTTGGAGGGAAGAGGCCTCGAAACTGAC 1620

SUBSTITUTE SHEET

FIG. 40.

27/42

BspMI
 |
 CCAAAGGGCAGGTCTCTATGCCCTTTCAGAACGGAGGAACGTGGCAGTTTGATTTCAA 1680

 HgiAI
 Bsp1286 EcoNI
 |
 CAGTGAGCACCTCTTAGCCTAAGCCAGGGCTGCTCAATTCTCTGGAGTCTCCTCGCTACC 1740

 EcoO109
 DraII
 Eco47III
 CelII HaeII
 | | |
 ATAAAAATGCTCAGCGCTGGGTCCCTGGGTTTGTGACTGACTCTGACTTCCCATGATGGCTT 1800

 StuI
 EaeI
 | |
 TTGCTCTGGAATGGACCCCTTCTCCTTAGTAACAGGCCCTCTTACCACAAATCTTCGTTTTT 1860

 EcoO109
 BspMI DraII HaeII
 | | |
 TTTTTTTAAATGCTGTTTTCAAAAAGTGAGAGGCAGGTCTCAACCAACCCCTGGAGCGCT 1920

 Bsp1286 NsiI
 |
 CCAGAAGCCCAGGCGTGCCCTCATGCAATTTCTCTGTGGTCTCTTGGCCCCAGACCTCCT 1980

FIG. 4p.

HgiAI
Bsp1286
|
GTTTGATTGGATAACTGCATTTTATATACTGAGCACGCTCTAAGTGGTCTTTATTTTAT 2040
TTTCCCTATCGAGTGTGTAGATGAAGAGTGATGACAATCCTGTAAATGTACTAGAACTT 2100
XmnI
|
TTTTATTAAAGGAACCTTTTCCCAAAAAAAAAAAAAAAAAAAAAA 2156

28/42

E-cadherin restriction map

BanI
|
CGGGCACCTGTGATTCCGGGAAGTCTCTGCCGCCCTCGCGCCGCTCGCGCCGCTCTCGA 60
HaeII
BbeI
NarI
AhaII
Bani
|
CCCCCGCCCGCCATGGGCCCTCGGTACGGGGCGGCCCGGCTCTGCTCCCCGCTGCTG 120
BspMI PstI BanII BglI
|
CTGCTGCTGCAGGTCTCATCGGGGCTCTGCCAAGAGCCGGAGCCCTGCGCCCTGGCTTT 180

FIG. 4q.

29/42

HaeII AflIII
|
GGCGCTGACAGCTACACGTTACCCGTGCCCCGGCGACACTTGGAGAGAGGCCGTGTCCTG 240

StyI
|
Cfr10I AvrII
| |
GGCAGGGTGAGTTTGAAGGATGCACCGGTCTACCTAGGACAGCCCTATGTTTCTGATGAC 300

AccI
|
ACCCGATTCAAAGTGGGCACAGATGGTGTGATTACAGTCAAGCGGCCCTCTACAACTTCAT 360

BspHI
|
AAACCAGAGATAAGTTTCTTGTCCATGCCCTGGGACTCCAGCCGCAGGAAGCTCTCCACC 420

HgiAI
|
AGAGTTAGGCTGAAGGCAGCGACGCCACCACCACCACCATCATGATGCTCCCTCTAAA 480

PvuII EaeI
| |
ACCCAGACAGAGGTGCTCACATTTCCAGTTCCTCCAGCATGGACTCAGAAGACAGAAGAGA 540

GACTGGGTTATCCCTCCTATCAGCTGCCCGGAAACGAGAAAGGCCCATTTCCCTAAAAC 600

BalI
|
CTGGTTCAGATCAAGTCTAACAGGGACAAAGAAATCAAGGTTTCTACAGCATCACTGGC 660

StyI

FIG. 4r.

30/42

720

CAAGGAGCTGACGCACCTCCTGTTGGTGTGTTATTATTGAAAGAGAAACAGGATGGCTG

780

AAGTGACTGAGCCTCTGGATAGAGAACAAATTGCTAAGTACATTCTCTACTCTCATGCC

BsmI

BclI

840

GTATCTTCTAATGGGAATGCGGTTGAAGACCCCAATGGAGATCGTGATCACGGTGACAGAT

AvaI

StyI

BamI

900

CAGAAATGACAACAAGCCCGAGTTCACCCAGGCAGTCTTCCAAGGATCTGTCTACGGAAGGT

BamI BspMI

XhoII

960

GCCCTTCCAGGCACCTCTGTGATGCAGGTGACAGCCACAGATGCGGATGATGTGAAT

1020

ACCTACAACGCTGCCATCGCTTACAGCATCCTCACACAAGACCCCTCCTGCCTAGCAGC

HgiAI

BstXI

1080

ATGATGTTCACTATCAACAAGGACACAGGAGTCATCAGCGTGCTCACCACTGGGCTGGAC

StyI

BspMI

1140

CGAGAGGGTGTCCCCATGTACACCTTGGTGGTTTCAAGGCTGTGACCTGCAAGCGGAAGGC

SUBSTITUTE SHEET

FIG. 4s.

```

      BclI      PvuII      AlwNI      |
      |         |         |
TTAACTACAAGTGAACAGCTGTGATCACAGTCACTGACATCAATGATAACCCCCCATC 1200

      BanI      |
      |         |
TTCAACCCAAACCAAGTACCAAGGACGGGTGCTGAGAACAAAGGCTAACGTCGAAATCGCT 1260

      BglI      |
      |         |
GTAATAAGTACGATGCTGATGTCCTCCGATACCCCGGCTGGAGGGCTGTGTACACC 1320

      BclI      |
      |         |
ATATTGAACAATAACAATGATCAATTGTTGTCAACCAAGACCCAGTAACTAACGACGGC 1380

      |         |
ATTTGAAAAACAATAAGGCTTGGATTTTGAGGACAAGCAGCAGTATGCTCTGTACGTG 1440

      AlwNI      |
      |         |
ACTGTGGTGAACGTGACCCCGTTTGAGGTCACTCCTCTCCACCTCCACAGCCACTGTCACT 1500

      |         |
GTGGACGTGGAAAGATGTGAATGAAGCCCCCATCTTCATCCCTTGCCCCAAAGGTAGTGTC 1560

      XhoII      BamHI      |
      |         |
ATCCCTGAAGACTTTGGTGTGGGCCAGGAAATCACATCCTACACCGCCGAGGATCCAGAT 1620

      Cfr10I      |
      |

```

SUBSTITUTE SHEET

32/42

FIG. 4t.

ACATATATGGAACAGAGGATAACGTATCGGATTGGAGGGATGCTGCCGTTGGCTGGAG 1680
 BamI
 PflMI
 AlwNI
 |
 GTTAATCCAGAATCTGGTGCCATTTTCACTCGGGCTGAGCTGGACAGAGAGGATTTTGAG 1740
 AvaI
 |
 CeuII
 |
 CACGTGAAGAAATAGCACGTATGAAGCCCTCATATAGCCATTGACTTCGGTTCTCCAGTT 1800
 GCTACTGGAACGGGAACTCTTCTACTGGTCCCTCTCTGATGTGAATGACAATGGCCCCATT 1860
 CCAGAACCTCGAAATATGGACTTCTGCCAGAAAACCCACAGCCTCATGTCAATCAACATC 1920
 XhoII
 BglII
 |
 ATTGATCCAGATCTTCCCCCAACACATCTCCCTTCACAGCAGAACTAACACACGGCGCA 1980
 HincII
 |
 AGTGTCAACTGGACCATCGAGTACAATGACCCAGCTCGTGAATCTCTAATTTTGAAGCCA 2040
 AAGAAAACCTTAGAGTTGGGTGACTACAAAATAAATCTCAAGCTCACAGATAACCAGAAC 2100
 BstEII
 |
 PvuII
 |
 HincII
 |
 AAGGACCAGGTGACCACCCCTATATGTGTTTGTGTGCGACTGCCGAAGGTGTCGTCAACAGC 2160

SUBSTITUTE SHEET

FIG. 4w.

35/42

DraI AhaIII	3120
ACTTGTCTCATTTTTTAAAGGAAGGTAGGGCTAAACTACCCCTATTGTGTTGTGTGT	
GTGTGTGTATGTGTAAATATTTTTTAATTGTGTCTCTTTTTTCTCCTATCACTGCACCTGGT	3180
EcoNI	3240
GTCCCGTGTCTAATAACCACTCTTAACCTCCTTCTGAACCTTACATTGCCCTCAGACAGGAG	
BanII ApaI EcoO109 DraII PstI EaeI	3300
TTCTCTGCTGCAGAAATTAATGGGCCCTTTCAGGATAAGAGACTTGGTCTTAGTTGATG	
Tth111I	3360
GTAGTGTGACTGGGTATTATGGACTCGTAAGGACTTTTAGTGGTCTCCTTTTATTATTTCC	
AatII AhaII	3420
TAGTACATAAAATTGAAATTCATATCCACTCCACTGACTTGTCTGCATTAAAGTGTGTTTG	
AatII AhaII	3480
TCATGTGGACGTCATTATTGGGCTACTTTTGGTTCTGAACAAGGAGCATTGACCAGAAAAG	
BanI	3540
GTGGTGAAATTTTCAGGTGCCACTCAACTCTTAATGTTCACCTATCACTCAACAGAAAGAG	

SUBSTITUTE SHEET

37/42

DrAI
AhaIII
|
GCAAAGGAAGGTGGGAGAGCCTTGACTTGATTTTTTTAAATTGAAATGTGAACTTC 4020

StyI
NcoI
|
AAGGAACTTTTGACAACCATGGGAAATAATTTATCTTAAATTGCTTTACTGTCTGTCTCAG 4080

PvuII
|
CTGTTTTTCAAAGAAAAAATCATCCCTGCAATCATTCTTGGAAATGTCTTGATTT 4140

DrAI
AhaIII
|
TTCAGCAATTAAACTCTAATTTAGTCCTGTATAGAGAAATGTTAATGTAGTTTTCAGTGT 4200

ATATGTGTGTGGGTACGGATAATTTTGTAATTTTCTTTAGCTCTGGAAAAGGAAACAATT 4260

SspI
|
TAAGCTGCCGAAAATTCTTAAATATTCATTTTTTATAAATTTTATTAAAGAAATTTTGTAA 4320


AAAAAAAAAAAAA 4333

FIG. 4y.

SUBSTITUTE SHEET


38/42

FIG. 5.



39/42

FIG. 6.



40/42




FIG. 7.

41/42




FIG. 8.

42/42




FIG. 9.

INTERNATIONAL SEARCH REPORT

International Application No PCT/US90/05105

I. CLASSIFICATION OF SUBJECT MATTER (if several classification symbols apply, indicate all) *

According to International Patent Classification (IPC) or to both National Classification and IPC

IPC(5): A61K 37/02, 39/00; C07K 7/08, 7/10, 13/00, 15/00, 15/28

U.S. Cl.: 530/324, 326, 350, 389, 390, 391, 402, 409, 345, 387; 514/12, 13; 424/85.8, 85.91

II. FIELDS SEARCHED

Minimum Documentation Searched *

Classification System :

Classification Symbols

530/324, 326, 350, 389, 390, 391, 402, 409, 345, 387
514/12, 13

U.S. Cl. 424/85.8, 85.91

Documentation Searched other than Minimum Documentation
to the Extent that such Documents are Included in the Fields Searched *

Data bases: Dialog (Files; Medline, Biosis, Chemical Abstracts, World Patents Index) Automated Patent Searching (1975-1990)

III. DOCUMENTS CONSIDERED TO BE RELEVANT ¹⁴

Category *	Citation of Document, ¹⁴ with indication, where appropriate, of the relevant passages ¹⁵	Relevant to Claim No. ¹⁶
<u>X</u> Y	The EMBO Journal, Volume 4, No. 13A, issued December 1985, Vestweber et al., "Identification of a Putative Cell Adhesion Domain of Uvomorulin," pp. 3393-3398. See the Abstract and Discussion.	1-6,14-21,23-27 & 35-42 1-65
Y	Development, Volume 102, issued April 1988, M. Takeichi, "The Cadherins: Cell-cell Adhesion Molecules controlling Animal Morphogenesis," pp. 639-655 see the Summary and pages 643, 645 and 651.	1-65
<u>X</u> Y	The Journal of Cell Biology, Volume 107, issued October 1988, B. Gumbiner et al., "The Role of the Cell Adhesion Molecule Uvomorulin in the Formation and Maintenance of the Epithelial Junctional Complex," pp. 1575-1587 see the Abstract.	1-6,14-21,23-27, 35-42 1-6,14-27,35-47, 55-65

* Special categories of cited documents: ¹⁵

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

IV. CERTIFICATION

Date of the Actual Completion of the International Search *

Date of Mailing of this International Search Report *

21 November 1990

04 FEB 1991

International Searching Authority *

Signature of Authorized Officer ¹⁷

R. Keith Baker
R. Keith Baker, Ph.D.

TSA/US

III. DOCUMENTS CONSIDERED TO BE RELEVANT (CONTINUED FROM THE SECOND SHEET)		
Category *	Citation of Document, ^{1a} with indication, where appropriate, of the relevant passages ^{1b}	Relevant to Claim No ^{1c}
Y	The ENBO Journal, Volume 6, No. 12, issued 1987, M. Ringwald et al., "The Structure of Cell Adhesion Molecule Uvomorulin Insights into the Molecular Mechanism of Ca ⁺⁺ -dependent Cell Adhesion," pp3347-3353, see pages 3647-3648.	1-13,22-34,43-54 and 63-65
Y	US, A, 4.671,958 (Rodwell et al.) 09 June 1987, see the Abstract and Column 7.	43-47 and 55-65
Y,P	Development Biology, Volume 139, No. 1, issued May 1990, O.W. Blaschuk et al., "Identification of a Cadherin Cell Adhesion Recognition Sequence," pp227-229, see the entire Document.	1-65

FURTHER INFORMATION CONTINUED FROM THE SECOND SHEET

V. ☐ OBSERVATIONS WHERE CERTAIN CLAIMS WERE FOUND UNSEARCHABLE¹

This international search report has not been established in respect of certain claims under Article 17(2) (a) for the following reasons:

1. ☐ Claim numbers _____, because they relate to subject matter not required to be searched by this Authority, namely:

2. ☐ Claim numbers _____, because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out¹, specifically:

3. ☐ Claim numbers _____, because they are dependent claims not drafted in accordance with the second and third sentences of PCT Rule 6.4(a).

VI. ☒ OBSERVATIONS WHERE UNITY OF INVENTION IS LACKING²

This International Searching Authority found multiple inventions in this international application as follows:

See Attachment

1. ☒ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims of the international application. **Telephone Practice**

2. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims of the international application for which fees were paid, specifically claims:

3. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claim numbers:

4. ☐ As all searchable claims could be searched without effort justifying an additional fee, the International Searching Authority did not invite payment of any additional fee.

Remark on Protest

- ☐ The additional search fees were accompanied by applicant's protest.
☐ No protest accompanied the payment of additional search fees.

Attachment To PCT/ISA/210
Observations Where Unity Of Invention Is Lacking

Group I, claims 1-13 and 22-34, drawn to a composition for opening tight junctions and a method of use, classified in classes 530 and 514, subclasses 324, 326, 350 and 12 and 13, respectively.

Group II, claims 14-21 - 35-42, drawn to antibodies for opening tight junctions and methods of use, classified in classes 530 and 424, subclasses 387 and 85.8, respectively.

Group III, claims 43-54 and 63-65, drawn to a conjugates of a drug and a cell adhesion inhibitor, classified in class 530, subclasses 402, 409, and 345.

Group IV, claims 55-62, drawn to a conjugate of a drug and an antibody, classified in classes 530 and 424, subclasses 389, 390, 391 and 85.91, respectively.

Attachment To PCT/ISA/210

Detailed Reasons For Holding Lack Of Unity Of Invention:

PCT Rule 13.2 permits claims to "a" (one) product, "a" (one) method of making and "a" (one) method of using said product. The invention as set forth in Group I constitutes a combination of a product and a method of use. Groups II, III and IV one drawn to products that are distinct from that of Group I. Each of the products have a different structure and are distinct compositions as evidenced by their separate classification.

ORIGINAL REVISION PAGE 2

THIS PAGE BLANK (USPTO)

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☒ **BLACK BORDERS**
- ☐ **IMAGE CUT OFF AT TOP, BOTTOM OR SIDES**
- ☐ **FADED TEXT OR DRAWING**
- ☐ **BLURRED OR ILLEGIBLE TEXT OR DRAWING**
- ☐ **SKEWED/SLANTED IMAGES**
- ☐ **COLOR OR BLACK AND WHITE PHOTOGRAPHS**
- ☐ **GRAY SCALE DOCUMENTS**
- ☐ **LINES OR MARKS ON ORIGINAL DOCUMENT**
- ☐ **REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY**
- ☐ **OTHER:** _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.

THIS PAGE BLANK (USPTO)